

# ***Retrieval of Biophysical/Structural Canopy Properties: An evolving synthesis of imaging spectrometry data and models***

J. Miller, P. Zarco-Tejada, D. Haboudane,  
W. Verhoef, S. Jacquemoud, A. Gitelson,  
I. Moorthy, H. Bach, D. Riaño,  
G. Schaepman-Strub, E. Chuvieco

*Canada, Spain, France, Netherlands, USA, Germany*

1. Earth Observations and Analysis Approaches - A perspective
2. Analysis Approach Selection: Rationale and Challenges
3. Analysis Paradigm Development and Assessment
4. Variable Retrieval Maps: Forestry, Agriculture, Landscape
5. New Developments
6. Suggested Next Steps

Imaging  
Spectrometer data:  
full spectrum, multi-view,  
multi-temporal

calibration,  
atmospheric correction,  
spectral polishing,  
etc.

-> TOC BRF

(thank you Alex et al.) !!!

Analysis Paradigms: Combinations of  
Canopy/leaf reflectance models  
Feature fitting based on physical/empirical models  
Optimized vegetation indices & RT models  
Linear/Non-linear SMA  
Spectral matching  
Artificial Neural networks trained with model runs  
Vegetation growth models

Environment & Resource  
Management “Products” of Interest

Agriculture:

e.g. Yield forecast map

Crop spatial/temporal growth variability maps

Crop stress map (nitrogen deficiency, insect, disease, dehydration, senescence)

Forestry:

e.g. Forest inventory map (e.g., forest area, forest type, fragmentation, stem volume)

Productivity map

Forest carbon map (reforestation, afforestation, deforestation)

Forest condition map (e.g., health, water stress, fuel type)

the current & most important challenge

Biophysical Variables:

LAI, Leaf Chlorophyll content

Leaf or Canopy water content

Bio-indicators: canopy chemistry, pigment ratios

Fractional cover, APAR, biomass

Spectral albedo

Clumping index


increased accuracy through remote sensing science innovations

## Implicit Assumptions

- hyperspectral and/or selected narrow band multispectral image data available at required time resolution
- appropriate/validated RT models across spatial scales of target pixel

## Potential Advantages Demonstrated

- yields quantitative estimates of target variables that are inherent surface properties, insensitive to location and time of observation
- ground truth ‘calibration’ not essential
- rapid delivery of product maps feasible
- responsive to worldwide scientific efforts to develop and improve appropriate RT models and techniques for variable(s) retrieval (i.e. can exploit remote sensing science advances)



“Comparison” can involve VERY different retrieval paradigms



### Forest Canopies



(Numbered in order of difficulty: -1 to 3 tackled -4, 5 underway)

#### Closed canopies

- Negligible background influence, only shadowing

#### Sparse canopies

-background influence throughout, with spectral distinctiveness of background critical

#### Clumped canopies

-background influence between crown, with spectral distinctiveness of background critical

### Agriculture Canopies





Numerous canopy models under development and intercomparisons for specific configurations are available

(a) - Side view



(b) - Top view



**Pinty et al., 2004. Radiation Transfer Model Intercomparison (RAMI) exercise**

*Specific canopy/biome/ecosystem conditions allow a judicious choice of RT model for quantitative variable retrieval – authors have experience with*

- SAILH, SAIL++, 4-SAIL, Kuusk-Nilson
- FLIM, GeoSAIL, rowSAIL, SPRINT, FLIGHT (each coupled with the PROSPECT leaf model)

*- chosen as appropriate to canopy closure, density, clumping, closure, background contrast, etc.*

**1. Utilize Airborne Hyperspectral Imagery Collection for Forestry and Precision Agriculture Tests**

- ~ 2 to 4 m spatial resolution hyperspectral imagery
  - ~ <1 m resolution with selected narrow bands nadir view & possible multi-angle imagery potential
  - multi-temporal (3 or 4 campaigns per season, varying SZA)
  - multi-year (2 or 3 successive years)
- to evaluate spatial heterogeneity
  - scale of relevance to “users” & measure background R
  - evaluate t/s robustness
  - evaluate stability


**2. Select (or manipulate) sites for a wide range of biophysical variable values**

**3. Field measurements by Collaborators (AAFC, OFRI) at each scale to Validate/Evaluate RT Models at leaf, canopy, airborne and satellite scales including:**

- top-of-canopy leaf sampling
- leaf biochemistry (pigments, water, etc) (AAFC, OFRI) , and optical r, t (York) for leaf RT model characterization/evaluation
- atmospheric characterization - AOD (Aeronet)
- structural, fCover & understory: LAI, LIDF, fCover,  $R_b(I)$ , height, clumping

**4. Generate maps of predicted canopy variables for assessment, evaluation, and use and feedback by collaborators to demonstrate robustness, accuracy, temporal/spatial stability**


*Evaluate effect of Model Assumptions and Real World Canopies*






Canopy **BRF** simulation

Leaf  $\rho$  &  $\tau$  simulation



Canopy **BRF** simulation

Leaf  $\rho$  &  $\tau$  simulation



## Algorithm Evaluation Completed or Underway:

### Uniform, extended, dense, closed canopies

- deciduous forests
- cereal crops near maturity

### Extended, thin, open canopies

- cereal crops shortly after emergence
- sparse poplar forest stands

account for  $\Delta$  background

### Open, dense, clumped canopies

- olive groves, vineyards
- conifer forests






Deciduous Sugar Maple closed canopy  
 - leaf chlorophyll retrieval (LAI > 3)  
 - widely separated individual sites  
 - canopy shadowing insensitive

Paradigm: inverse RT models  
 (PROSPECT-SAILH)  
 with VI (R750/R710) as “best”  
 merit function for Cab

Application: permits Cab mapping into  
 3 to 4 bio-indicator “stress” levels,  
 given known species maps





Leaf  $Chl_{a+b}$  by numerical model Inversion (SAILH + PROSPECT) for closed canopies of sugar maple.




Low stressed site

Measured: 38.8  $\mu\text{g}/\text{cm}^2$   
 Estimated: 35.2  $\mu\text{g}/\text{cm}^2$

High stressed site

Measured: 19.1  $\mu\text{g}/\text{cm}^2$   
 Estimated: 20.2  $\mu\text{g}/\text{cm}^2$






RMSE (and  $R^2$ ) for LAI prediction were 0.28 (0.98), 0.46 (0.89), and 0.85 (0.74) for soybean, corn and wheat respectively


Paradigm: Up-scaling new VI: MTVI2 based on PROSPECT-SAILH simulations

Application: Seasonal progression of LAI development (< 1 week turnaround) for input into productivity estimates.







-Note sig. difference in r, t of broadleaves




Analysis Paradigm:  
inverse RT models (PROSPECT-SAILH) for isolated tree crowns with VI (R750/R710) as merit function for Cab











JUNE



SEPTEMBER






Olive groves in Andalucia, Spain




Extension of modeling approach to clumped canopies with soil understory - underway  
(Zarco-Tejada et al , RSE & Brugges, 2004)

Vineyards in Castilla-Leon, Spain







Olive Grove Study sites




Vineyard Study sites








Analysis Paradigm: Cab estimation from crowns only using SAILH-PROSPECT simulation for MCARI/OSAVI VI prediction

Note: For applicability to coarser spatial resolution sensors, clumped canopy predictions using FLIM and SPRINT also investigated



Application: Crown chlorophyll pigment image showing spatial patterns for precision agriculture management action or potential segregation by oil quality







Analysis Paradigm: Cab estimation at canopy level by scaling up TCARI/OSAVI through PROSPECT + rowMCRM model

Application: Cab variation as a potential surrogate for grapes segregation for wine quality


1. Leaf/canopy model




2. Training ANN




3. Application to image





Analysis Paradigm: Artificial Neural Network (ANN) trained with canopy reflectance spectra simulations with i) PROSPECT-SAILH, ii) PROSPECT-rowMCRM, and iii) PROSPECT-FLIM models, for grassland, row crop, forest cover types, respectively, for increments in Leaf EWT and canopy LAI.







1. Earth Observations and Analysis Approaches - A perspective
2. Analysis Approach Selection: Rationale and Challenges
3. Analysis Paradigm Development and Assessment
4. Variable Retrieval Maps: Forestry, Agriculture, Landscape
- 5. New Developments**
6. Suggested Next Steps

Analysis Paradigm: SLC (Soil-Leaf-Canopy) model [4SAIL2 canopy model, PROSPECT leaf model, Hapke Soil model, vertical leaf greenness gradient as in GeoSAIL, clumping as in FLIM] for simulation of surface bi-directional reflectances as seen with CHRIS from the PROBA satellite. Inversions with LUT on a pixel-by-pixel comparison of observed and modeled canopy reflectances with minimization of RMS error on full spectra (Bach et al., 2005).

Applications: Added retrieval of structural parameter LADF, expected to provide improved accuracy in LAI retrievals.



*Simulated relative-BRDF factor at 773nm a maize field with LAI of 3.5 and its change with the canopy structure parameter LIDF (top), compared with two BRDF factors for maize measured by CHRIS on July 18 and August 03 (right)*



*Retrieval results using SLC and five angular CHRIS acquisitions on Jul 18;  
left: LIDF-a equivalent to average leaf angle; right: Leaf Area Index*

Bach & Verhoef (2006)



F  
I  
G  
O  
S



C  
H  
R  
I  
S



A  
S  
A  
S  
P



M  
O  
D  
I  
S

Inst. multi-angular  
high angular resolution

Inst. multi-angular  
low angular resol.

Daily composites  
by flight patterns

Multiple-day  
composites

Numerical  
integration

Atmospheric correction (aerosols, surface anisotropy, diffuse illumination)  
BRDF model inversion (e.g., MRPV, Roujean, Ross/Li)  
$$BRF_{\lambda}(\theta_i, \theta_r, \Phi_{i-r}) = K_{iso, \lambda} + a_{\lambda} K_{geo}(\theta_i, \theta_r, \Phi_{i-r}) + b_{\lambda} K_{vol}(\theta_i, \theta_r, \Phi_{i-r})$$

BRDF forward modeling for any ratio of direct to diffuse illumination

$BHR_{\lambda}$

$BHR_{\lambda}$  - albedo;  $BHR_{iso, \lambda}$  - white-sky albedo;  $DHR_{\lambda}$  - black-sky albedo





Ambient  
conditions

Few studies  
Atm. correction

No albedo products  
Co-registration

Few studies  
Atm. correction

Spectral resol.  
Surface changes



Spectral albedo (BHR) based on field goniometer measurements (numerical integration)



Spectral albedo (DHR) based on HyMap multi-angular flight pattern data and BRDF model

Isotropic kernel, geometric and volumetric kernel weights

-> spectral dependence!


Courtesy: U. Beisl, 2001





## ***Status and Recommendations for Spectral Albedo Derivation based on Imaging Spectroscopy***

- Spectral land surface albedo has a high potential for more detailed representation of vegetation dynamics in climate modeling, is used for validation purposes of broadband satellite albedo products and in meteorological studies.
- No current spectral albedo product based on high spectral resolution (>7) and instantaneous multi-view-angular satellite data.
- Satellite, airborne and in-situ based spectrometer data have to be corrected for the diffuse irradiance!  
Neglecting the diffuse irradiance results in distortions of the derived BRDF, and thus in spectral albedo uncertainties (especially in blue spectral wavelengths).
- Spectral surface albedo databases have to be standardized in terms of description of the provided physical quantity, vegetation (e.g., LAI, LAD, stem density), solar angle, and atmospheric conditions.



Temporal behavior of maize biophysical characteristics from detailed field data - A. Gitelson





- Through model-based approaches repetitive hyperspectral images over vegetated targets can predict canopy-leaf variables (specifically Leaf Cab, LAI, Leaf EWT, fCover) with sufficient accuracy to track seasonal, spatial, species-dependent changes measured in the field
- Maps of such variables have been shown to provide valuable information in forestry and precision agriculture applications, often indirectly related but correlated to desired products: inventory, speciation, land cover, stress (condition) maps, productivity/yield prediction/potential maps
- Multi-temporal retrieved variable maps have significant potential for model-based product retrieval through crop growth models: eg. crop biomass & yield prediction

- Increase emphasis on multi-angle imaging spectroscopy data because of new variables for retrieval (eg. spectral albedo, LIDF), and increased retrieval accuracy for primary biophysical variables
- assess new vegetation indices related to crop growth parameters (APAR, fCover, biomass) through leaf-canopy model simulations and refine such VIs for use in rapid variable map production through scaling up
- Improvements to PROSPECT to (i) provide Ca/Cb ratios, and (ii) better represent needle leaves, would open a significant new range of applications and potential
- Exploit and continue development of FluorMOD (a version of PROSPECT and 4SAIL which simulates TOC and TOA reflected radiances as well as fluorescence emissions) due to (i) potential to affect observed radiance in red and NIR regions and thus affect retrievals, and (ii) potential complimentary information – vegetation photosynthetic functioning



*Thank you for your attention!*




**Paradigms for deriving bio- geophysical variables from above-canopy bi-directional reflectance :**

**I. Canopy-Leaf model inversions: **biophysical variables** are retrieved by**

- (i) inverting RT models and matching complete spectra in the merit function (e.g Bach & Verhoef, 2006)
- (ii) inverting RT models using key optical indices in the merit function for inversion (eg. Zarco-Tejada et al (2001) for deciduous closed canopy forests; Moorthy et al (2003) for boreal conifer forests; Zarco-Tejada et al (2004) for olive groves, vineyards)



Merit Function

$$\Delta^2 = \sum_{i=1}^n [r_{obs}(\lambda_i) - r_{mod}(\lambda_i, P)]^2$$




**P**= set of parameters: **Cab, LAI, θL...**

**II. Upscaling through RT models simulations: **new optical indices** (eg. TCARI/OSAVI, MTVI, etc) sensitive to **specific biophysical variables** (with minimal sensitivity to other variables) are **simulated at above-canopy level** (e.g. Haboudane et al 2002, 2003, for agriculture crops) forming the basis of a retrieval algorithm**








**C<sub>ab</sub> estimation from ROSIS crowns using MCARI/OSAVI prediction**



**C<sub>ab</sub> estimation from aggregated pixels (C+Sh+So) using MCARI/OSAVI**





**Crowns**  
**PROSPECT-SAILH**




**Crown+Shadow+Soil**  
**PROSPECT-SAILH &**  
**PROSPECT-SAILH-FLIM**


Maps of Crop Fraction  
Estimated:  
- (i) Using Linear Spectral  
Unmixing  
- (ii) Using VIs: TSAVI,  
OSAVI, MTVI2

found comparable in  
performance, but with  
advantages &  
disadvantages in  
implementation








Soil Brightness and Colour



Soil Principal Components



1. Earth Observations and Analysis Approaches - A perspective
2. Analysis Approach Selected: Rationale and Challenges
3. York Algorithm Development Strategy and Steps
4. Variable Retrievals: Forestry & Agriculture 1<sup>st</sup> Steps
- 5. Towards Products: Forestry & Agriculture, Next Steps**
6. Satellite HIS Issues
7. Conclusions



## Products for Agriculture & Forestry (eg)


- cover map – inventory, speciation, land cover
- stress (condition) map – water stress, nitrogen stress, etc
- productivity/ yield prediction/potential map
- carbon map

## Applicability of Model-based approach?

Variables can in many cases be directly retrieved, with sufficient accuracy and flexibility.

Products not directly retrievable – require new approaches & developments to exploit model-based advantages

--- some examples



IFC -2: LAI

IFC -2: Dry Shoot Biomass

Ground Measured Yield

# Derived Leaf Pigment Content, as a surrogate for Land Cover?


IGARSS 2006 & 27<sup>th</sup> CSRS



Boreal Forest near Sudbury, Ontario Canada 2.2 x 1.5 km

## Land Cover Issues:

Patterns of estimated leaf pigment content over boreal forests (north Ontario) are indicative of land cover patterns



1. Earth Observations and Analysis Approaches - A perspective
2. Analysis Approach Selected: Rationale and Challenges
3. York Algorithm Development Strategy and Steps
4. Variable Retrievals: Forestry & Agriculture 1<sup>st</sup> Steps
5. Towards Products: Forestry & Agriculture, Next Steps
- 6. Satellite HIS Issues**
7. Conclusions



## Issues

- required spatial resolution for specific applications
- challenges to the model-based approach for variable retrieval

Methodology: Haboudane *et al.*, 2002)

⇒ **CARI** → minimize effects of non-photosynthetic materials

⇒ **MCARI** → depth of chlorophyll absorption at 670 nm relative to the reflectance at 550 nm and 700 nm  
still sensitive to non-photosynthetic element effects

$$MCARI = [(R_{700} - R_{670}) - 0.2 \cdot (R_{700} - R_{550})] \cdot \left( \frac{R_{700}}{R_{670}} \right)$$



⇒ **TCARI** → improving its sensitivity at low chlorophyll values

$$TCARI = 3 \cdot [(R_{700} - R_{670}) - 0.2 \cdot (R_{700} - R_{550})] \cdot \left( \frac{R_{700}}{R_{670}} \right)$$




⇒ Influence by soil reflectance for low values of LAI  
→ **OSAVI** minimizes soil effects with MCARI

$$OSAVI = (1 + 0.16) \cdot \left[ \frac{R_{800} - R_{670}}{R_{800} + R_{670} + 0.16} \right]$$

**TCARI/OSAVI**



**Cornfield at L'Acadie Experimental farm, Agriculture  
and Agri-Food Canada**



Paradigm: Up-scaling of TCARI/OSAVI leaf pigment algorithm derived from PROSPECT-SAILH simulations

Application: Spatial heterogeneity mapping for N application in precision farming

Haboudane et al RSE (2002)

## **Status and Recommendations for new VI algorithms for biophysical variables for crops**

- New VI Algorithms for the remote estimation of fraction of Absorbed Photosynthetically Active Radiation, Vegetation Fraction, Green LAI and Green Leaf Biomass were developed and successfully applied in **maize** and **soybean** crops
- Validation using independent datasets proved the suitability of the algorithms to monitor crop development along the growing season, as well as inter-annual variability
- As these algorithms can be viewed as empirical models, future work should focus on comparing leaf-canopy model simulations of the the VI algorithm performances in order to infer theoretical limits or provide improvements in formulation